arXiv:math/0401099v1 [math.GM] 10 Jan 2004

Elementary evaluation of $\int_0^\infty \frac{\sin^p t}{t^q}\mathrm{d}t$

Omran Kouba
Department of Mathematics
Higher Institute for Applied Sciences and Technology
P.O. Box 31983, Damascus, Syria.
E-mail : omran_kouba@hiast.edu.sy

Abstract

Let p and q be two positive integers. The goal of this note is to demonstrate, in a very simple and elementary way and without using advanced tools, a formula to express the value of the integral $I(p,q) = \int_0^\infty \frac{\sin^p t}{t^q}\mathrm{d}t$ when it converges.

Introduction :

Consider the sequence of polynomials $(P_n)_{n\geq 0}$ defined as follows :

$$P_{2n}(X) = \sum_{k=0}^{n} \frac{(-1)^k}{(2k)!} X^{2k}$$

$$P_{2n+1}(X) = \sum_{k=0}^{n} \frac{(-1)^k}{(2k+1)!} X^{2k+1}$$

Then Define, with the convention $P_{-1} = 0$, the following sequence of functions $(f_n)_{n\geq 1}$:

$$\forall n \geq 0, \qquad f_{2n+1} : \mathbb{R} \longrightarrow \mathbb{R}, f_{2n+1}(t) = (-1)^n(\sin t - P_{2n-1}(t))$$

$$f_{2n+2} : \mathbb{R} \longrightarrow \mathbb{R}, f_{2n+2}(t) = (-1)^{n+1}(\cos t - P_{2n}(t))$$

Our result concerning the evaluation of the integral $I(p,q)$ is based upon the next proposition which summarizes some properties of the functions $(f_n)_{n\geq 1}$:

Proposition 1. *The sequence of functions $(f_n)_{n\geq 1}$ satisfies the following properties :*

1°. *For every $n \in \mathbb{N}^*$ we have $f'_{n+1} = f_n$.*

2°. *For every $n \in \mathbb{N}^*$ we have $\lim_{t\to 0} \frac{f_n(t)}{t^n} = \frac{1}{n!}$.*

3°. *For every $n \in \mathbb{N}^*$ the improper integral $J_n = \int_0^\infty \frac{f_n(t)}{t^n}\mathrm{d}t$ converges, and*

$$J_n = \frac{1}{(n-1)!} J_1 = \frac{1}{(n-1)!} \cdot \frac{\pi}{2}$$

4°. *For every $n \geq 2$ the improper integral $K_n = \int_1^\infty \frac{f_{n-1}(t)}{t^n}\mathrm{d}t$ converges, and for all $\lambda > 0$ we have :*

$$\lim_{X\to\infty} \int_X^{\lambda X} \frac{f_{n+1}(t)}{t^n}\mathrm{d}t = \frac{\ln \lambda}{(n-1)!} \quad \textit{and} \quad \lim_{X\to\infty} \int_1^X \frac{f_{n+1}(t)}{t^n}\mathrm{d}t = +\infty$$

5°. *For every $(m,q) \in \mathbb{N}^2$ such that $1 \leq q \leq m$ we have*

$$\forall t \in \mathbb{R}, \qquad (\sin t)^{2m} = \frac{1}{2^{2m-1}} \sum_{k=1}^{m} \binom{2m}{m-k} (-1)^{k+q} f_{2q}(2kt) \tag{1}$$

6°. *For every $(m,q) \in \mathbb{N}^2$ such that $0 \leq q \leq m$ we have*

$$\forall t \in \mathbb{R}, \qquad (\sin t)^{2m+1} = \frac{1}{2^{2m}} \sum_{k=0}^{m} \binom{2m+1}{m-k} (-1)^{k+q} f_{2q+1}((2k+1)t) \tag{2}$$

PROOF. 1°. The verification of the first property is direct.

2°. The second property follows from

$$\cos t = P_{2n}(t) + \sum_{k=n+1}^{\infty} \frac{(-1)^k}{(2k)!} t^{2k} \quad \text{and} \quad \sin t = P_{2n-1}(t) + \sum_{k=n}^{\infty} \frac{(-1)^k}{(2k+1)!} t^{2k+1}$$

which is valid for all $n \in \mathbb{N}$ and all $t \in \mathbb{R}$.

3°. It is well known that the integral $J_1 = \int_0^\infty \frac{\sin t}{t} \mathrm{d}t$ converges and that $J_1 = \frac{\pi}{2}$. In the case $n \geq 2$ we have $\frac{|f_n(t)|}{t^n} = O\left(\frac{1}{t^2}\right)$ and this, with 2°, proves the convergence of the integral J_n. Now, for $n \geq 2$ and $X > 0$ we have

$$\begin{aligned} \int_0^X \frac{f_n(t)}{t^n} \mathrm{d}t &= -\frac{f_n(t)}{(n-1)t^{n-1}}\Big]_0^X + \frac{1}{n-1} \int_0^X \frac{f_n'(t)}{t^{n-1}} \mathrm{d}t \\ &= \frac{1}{n-1} \int_0^X \frac{f_{n-1}(t)}{t^{n-1}} \mathrm{d}t + O\left(\frac{1}{X}\right) \end{aligned}$$

Letting $X \longrightarrow +\infty$ we find that $J_n = \frac{1}{n-1} J_{n-1}$ and this proves 3° by induction.

4°. Considering two cases according to the parity of n we see immediately that :

$$\forall t > 0, \qquad \frac{f_{n+1}(t)}{t^n} = \frac{1}{(n-1)!} \cdot \frac{1}{t} - \frac{f_{n-1}(t)}{t^n} \tag{3}$$

But, for $n \geq 2$, we have $f_{n-1}(t) = O\left(t^{\max(n-3,0)}\right)$ in the neighborhood of $+\infty$, so that $\frac{|f_{n-1}(t)|}{t^n} = O\left(\frac{1}{t^2}\right)$, and this proves the convergence of $K_n = \int_1^\infty \frac{f_{n-1}(t)}{t^n} \mathrm{d}t$. Now, using (3), we conclude that $\lim_{X \to \infty} \int_1^X \frac{f_{n+1}(t)}{t^n} \mathrm{d}t = +\infty$ and that

$$\int_X^{\lambda X} \frac{f_{n+1}(t)}{t^n} = \frac{\ln \lambda}{(n-1)!} - \int_X^{\lambda X} \frac{f_{n-1}(t)}{t^n}$$

The convergence of K_n proves that $\lim_{X \to \infty} \int_X^{\lambda X} \frac{f_{n-1}(t)}{t^n} = 0$ so that

$$\lim_{X \to \infty} \int_X^{\lambda X} \frac{f_{n+1}(t)}{t^n} = \frac{\ln \lambda}{(n-1)!}$$

5°. Let us start with a well known standard calculation. Starting from Euler's formula : $\sin t = \frac{e^{\mathrm{i}t} - e^{-\mathrm{i}t}}{2\mathrm{i}}$ and using the binomial theorem we can write :

$$\begin{aligned} (\sin t)^{2m} &= \frac{(-1)^m}{2^{2m}} \left(\sum_{k=0}^{2m} \binom{2m}{k} (-1)^k e^{\mathrm{i}kt} e^{-\mathrm{i}(2m-k)t} \right) = \frac{(-1)^m}{2^{2m}} \left(\sum_{k=0}^{2m} \binom{2m}{k} (-1)^k e^{2\mathrm{i}(k-m)t} \right) \\ &= \frac{(-1)^m}{2^{2m}} \left(\sum_{k=0}^{m-1} \binom{2m}{k} (-1)^k e^{2\mathrm{i}(k-m)t} + (-1)^m \binom{2m}{m} + \sum_{k=m+1}^{2m} \binom{2m}{k} (-1)^k e^{2\mathrm{i}(k-m)t} \right) \\ &= \frac{(-1)^m}{2^{2m}} \left(\sum_{k=0}^{m-1} \binom{2m}{k} (-1)^k e^{2\mathrm{i}(k-m)t} + (-1)^m \binom{2m}{m} + \sum_{k=0}^{m-1} \binom{2m}{k} (-1)^k e^{-2\mathrm{i}(k-m)t} \right) \\ &= \frac{1}{2^{2m}} \left(\binom{2m}{m} + 2 \sum_{k=0}^{m-1} \binom{2m}{k} (-1)^{m-k} \cos(2(k-m)t) \right) \\ &= \frac{\binom{2m}{m}}{2^{2m}} + \frac{1}{2^{2m-1}} \sum_{k=1}^{m} \binom{2m}{m-k} (-1)^k \cos(2kt) \end{aligned}$$

Using that $\forall u \in \mathbb{R}$, $\cos u = (-1)^q f_{2q}(u) + P_{2q-2}(u)$, we can write

$$\forall t \in \mathbb{R}, \qquad (\sin t)^{2m} = Q^0_{m,q}(t) + \frac{1}{2^{2m-1}} \sum_{k=1}^{m} \binom{2m}{m-k} (-1)^{k+q} f_{2q}(2kt)$$

with $Q^0_{m,q}(t) = \dfrac{\binom{2m}{m}}{2^{2m}} + \dfrac{1}{2^{2m-1}} \displaystyle\sum_{k=1}^{m} \binom{2m}{m-k} (-1)^k P_{2q-2}(2kt)$.

Since, in a neighborhood of 0, we have, $\forall k \in \{1, 2, \ldots, m\}$, $f_{2q}(2kt) = O(t^{2q})$ and $(\sin t)^{2m} = O(t^{2m}) = O(t^{2q})$ (this results from the assumption $q \leq m$), we conclude that the polynomial $Q^0_{m,q}$, which is of degree at most $2q-2$, satisfies : $Q^0_{m,q}(t) = O(t^{2q})$ in a neighborhood of 0. This proves clearly that $Q^0_{m,q} = 0$, and establishes (1).

6°. In a similar way, starting from Euler's formula and using the binomial theorem we can write :

$$\begin{aligned}
(\sin t)^{2m+1} =& \frac{(-1)^{m+1}}{2^{2m+1}\mathrm{i}} \left(\sum_{k=0}^{2m+1} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}kt} e^{-\mathrm{i}(2m+1-k)t} \right) \\
=& \frac{(-1)^{m+1}}{2^{2m+1}\mathrm{i}} \left(\sum_{k=0}^{2m+1} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}(2(k-m)-1)t} \right) \\
=& \frac{(-1)^{m+1}}{2^{2m+1}\mathrm{i}} \left(\sum_{k=0}^{m} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}(2(k-m)-1)t} + \sum_{k=m+1}^{2m+1} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}(2(k-m)-1)t} \right) \\
=& \frac{(-1)^{m+1}}{2^{2m+1}\mathrm{i}} \left(\sum_{k=0}^{m} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}(2(k-m)-1)t} - \sum_{k=0}^{m} \binom{2m+1}{k} (-1)^k e^{\mathrm{i}(2(m-k)+1)t} \right) \\
=& \frac{(-1)^{m}}{2^{2m}} \sum_{k=0}^{m} \binom{2m+1}{k} (-1)^k \sin((2(m-k)+1)t) \\
=& \frac{1}{2^{2m}} \sum_{k=0}^{m} \binom{2m+1}{m-k} (-1)^k \sin((2k+1)t)
\end{aligned}$$

Using that $\forall u \in \mathbb{R}$, $\sin u = (-1)^q f_{2q+1}(u) + P_{2q-1}(u)$, we can write

$$\forall t \in \mathbb{R}, \qquad (\sin t)^{2m+1} = Q^1_{m,q}(t) + \frac{1}{2^{2m}} \sum_{k=0}^{m} \binom{2m+1}{m-k} (-1)^{k+q} f_{2q+1}((2k+1)t)$$

with $Q^1_{m,q}(t) = \dfrac{1}{2^{2m}} \displaystyle\sum_{k=0}^{m} \binom{2m+1}{m-k} (-1)^k P_{2q-1}((2k+1)t)$.

Since, in a neighborhood of 0, we have, $\forall k \in \{1, \ldots, m\}$, $f_{2q+1}((2k+1)t) = O(t^{2q+1})$ and $(\sin t)^{2m+1} = O(t^{2m+1}) = O(t^{2q+1})$ (this results from the assumption $q \leq m$), we conclude that the polynomial $Q^1_{m,q}$, which is of degree at most $2q-1$, satisfies : $Q^1_{m,q}(t) = O(t^{2q+1})$ in a neighborhood of 0. This proves that $Q^1_{m,q} = 0$, and establishes (2). □

The main result :

Proposition 2. *Consider* $(q, m) \in \mathbb{N}$.

1°. *If* $1 \leq q \leq m$ *then* $\displaystyle\int_0^\infty \frac{\sin^{2m}}{t^{2q}} \mathrm{d}t = \frac{\pi}{2^{2m}} \sum_{k=1}^{m} (-1)^{k+q} \binom{2m}{m-k} \frac{(2k)^{2q-1}}{(2q-1)!}$.

2°. *If* $0 \leq q \leq m$ *then* $\displaystyle\int_0^\infty \frac{\sin^{2m+1}}{t^{2q+1}} \mathrm{d}t = \frac{\pi}{2^{2m+1}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} \frac{(2k+1)^{2q}}{(2q)!}$.

3°. *If* $2 \le q \le m$ *then* $\displaystyle\int_0^\infty \frac{\sin^{2m}}{t^{2q-1}}\mathrm{d}t = \frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}\frac{(2k)^{2q-2}}{(2q-2)!}\ln k.$

4°. *If* $1 \le q \le m$ *then* $\displaystyle\int_0^\infty \frac{\sin^{2m+1}}{t^{2q}}\mathrm{d}t = \frac{1}{2^{2m}}\sum_{k=0}^{m}(-1)^{k+q}\binom{2m+1}{m-k}\frac{(2k+1)^{2q-1}}{(2q-1)!}\ln(2k+1).$

PROOF. 1°. Using formula (1) from Proposition 1.5°, we conclude that

$$\begin{aligned}\int_0^\infty \frac{\sin^{2m} t}{t^{2q}}\mathrm{d}t =& \frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}\int_0^\infty \frac{f_{2q}(2kt)}{t^{2q}}\mathrm{d}t\\ =&\frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}(2k)^{2q-1}\int_0^\infty \frac{f_{2q}(u)}{u^{2q}}\mathrm{d}u\\ =&\frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}(2k)^{2q-1}J_{2q}\end{aligned}$$

Now, using Proposition 1.3°, we conclude that

$$\int_0^\infty \frac{\sin^{2m} t}{t^{2q}}\mathrm{d}t = \frac{\pi}{2^{2m}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}\frac{(2k)^{2q-1}}{(2q-1)!}$$

2°. Using formula (2) from Proposition 1.6°, we conclude that

$$\begin{aligned}\int_0^\infty \frac{\sin^{2m+1} t}{t^{2q+1}}\mathrm{d}t =& \frac{1}{2^{2m}}\sum_{k=0}^{m}(-1)^{k+q}\binom{2m+1}{m-k}\int_0^\infty \frac{f_{2q+1}((2k+1)t)}{t^{2q+1}}\mathrm{d}t\\ =&\frac{1}{2^{2m}}\sum_{k=0}^{m}(-1)^{k+q}\binom{2m+1}{m-k}(2k+1)^{2q}\int_0^\infty \frac{f_{2q+1}(u)}{u^{2q+1}}\mathrm{d}u\\ =&\frac{1}{2^{2m}}\sum_{k=0}^{m}(-1)^{k+q}\binom{2m+1}{m-k}(2k+1)^{2q}J_{2q+1}\end{aligned}$$

Now, using Proposition 1.3°, we conclude that

$$\int_0^\infty \frac{\sin^{2m+1} t}{t^{2q+1}}\mathrm{d}t = \frac{\pi}{2^{2m+1}}\sum_{k=0}^{m}(-1)^{k+q}\binom{2m+1}{m-k}\frac{(2k+1)^{2q}}{(2q)!}$$

3°. Using formula (1) from Proposition 1.5°, we conclude that, for $X > 0$ we have :

$$\begin{aligned}\int_0^X \frac{\sin^{2m} t}{t^{2q-1}}\mathrm{d}t =& \frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}\int_0^X \frac{f_{2q}(2kt)}{t^{2q-1}}\mathrm{d}t\\ =&\frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}(2k)^{2q-2}\int_0^{2kX} \frac{f_{2q}(u)}{u^{2q-1}}\mathrm{d}u\\ =&\frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}(2k)^{2q-2}\int_{2X}^{2kX} \frac{f_{2q}(u)}{u^{2q-1}}\mathrm{d}u + C^0_{m,q}\int_0^{2X}\frac{f_{2q}(u)}{u^{2q-1}}\mathrm{d}u\end{aligned}$$

with $C^0_{m,q} = \displaystyle\frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}(2k)^{2q-2}$. But, for $2 \le q \le m$ the integral $\displaystyle\int_0^\infty \frac{\sin^{2m} t}{t^{2q-1}}\mathrm{d}t$ converges, so that using Proposition 1.4°, we conclude that we must have $C^0_{m,q} = 0$, and consequently

$$\int_0^\infty \frac{\sin^{2m} t}{t^{2q-1}}\mathrm{d}t = \frac{1}{2^{2m-1}}\sum_{k=1}^{m}(-1)^{k+q}\binom{2m}{m-k}\frac{(2k)^{2q-2}}{(2q-2)!}\ln k$$

4°. Using formula (2) from Proposition 1.6°, we conclude that, for $X > 0$ we have :

$$\begin{aligned}
\int_0^X \frac{\sin^{2m+1} t}{t^{2q}} \mathrm{d}t =& \frac{1}{2^{2m}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} \int_0^X \frac{f_{2q+1}((2k+1)t)}{t^{2q}} \mathrm{d}t \\
=& \frac{1}{2^{2m}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} (2k+1)^{2q-1} \int_0^{(2k+1)X} \frac{f_{2q+1}(u)}{u^{2q}} \mathrm{d}u \\
=& \frac{1}{2^{2m}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} (2k+1)^{2q-1} \int_X^{(2k+1)X} \frac{f_{2q+1}(u)}{u^{2q}} \mathrm{d}u + C_{m,q}^1 \int_0^X \frac{f_{2q+1}(u)}{u^{2q}} \mathrm{d}u
\end{aligned}$$

with $C_{m,q}^1 = \frac{1}{2^{2m}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} (2k+1)^{2q-1}$. But, for $1 \leq q \leq m$ the integral $\int_0^\infty \frac{\sin^{2m+1} t}{t^{2q}} \mathrm{d}t$ converges, so that using Proposition 1.4°, we conclude that we must have $C_{m,q}^1 = 0$, and consequently

$$\int_0^\infty \frac{\sin^{2m+1} t}{t^{2q}} \mathrm{d}t = \frac{1}{2^{2m}} \sum_{k=0}^{m} (-1)^{k+q} \binom{2m+1}{m-k} \frac{(2k+1)^{2q-1}}{(2q-1)!} \ln(2k+1)$$

And this achieves the proof. □

Numerical Values :

In the following table we find the values of $I(p,q) = \int_0^\infty \frac{\sin^p t}{t^q} \mathrm{d}t$ when this integral converges, calculated using Proposition 2.:

	$p = 1$	2	3	4	5
$q = 1$	$\frac{\pi}{2}$	–	$\frac{\pi}{4}$	–	$\frac{3\pi}{16}$
2	–	$\frac{\pi}{2}$	$\frac{3}{4} \ln 3$	$\frac{\pi}{4}$	$\frac{15}{16} \ln 3 - \frac{5}{16} \ln 5$
3	–	–	$\frac{3\pi}{8}$	$\ln 2$	$\frac{5\pi}{32}$
4	–	–	–	$\frac{\pi}{3}$	$\frac{125}{96} \ln 5 - \frac{45}{32} \ln 3$
5	–	–	–	–	$\frac{115\pi}{384}$
